

## FACTS AND FIGURES

Giorgos Mallinis, Ioannis Mitsopoulos, Petros Tsioras, Thomas Papachristou and Gavriil Spyroglou

### Definitions

Forests that resprout after felling

*πρεμνοφυή δάση (premnofie dasi)*

- Papachristou

Coppice forest, or paravlastogenes forest, is forest where regeneration is done by sprouts.

*Πρεμνοφυές ή παραβλαστογενές δάσος είναι το δάσος στο οποίο η αναγέννηση γίνεται με παραβλαστήματα.*

- Mallinis

### Legal Framework

Presidential Decree. 19-11-1928, 28-29.

Restrictions and guidelines regarding coppice forest harvesting.

### Statistics

Coppice forests cover an area of approximately 1,930,000 ha (12% of the total country's area). The main species managed as coppice are broad-leaved oaks (1,105,339 ha), beech (337,000 ha), chestnut (33,000 ha) and other broadleaved species (88,000 ha). The management of these coppice forests is intensive, with a clear cutting cycle ranging from 20 to 30 years.

1.1. Area of forest and other wooded land and its changes

	1964		1992	
	Area (1000 ha)	Percentage (%)	Area (1000 ha)	Percentage (%)
Forest *	2 512	19.0%	3 359	25.5%
Other wooded land *	3 960	30.0%	3 154	23.9%
<b>Forest and other wooded land</b>	<b>6 472</b>	<b>49.0%</b>	<b>6 513</b>	<b>49.4%</b>
Other land uses	6 724	51.0%	6 683	50.6%
<b>Total area</b>	<b>13 196</b>	<b>100.0%</b>	<b>13 196</b>	<b>100.0%</b>

\* Definitions are quoted in the Appendix I

1.1.1. Area of forest according to management type and its changes

Management type	1964		1992	
	Area (1000 ha)	Percentage (%)	Area (1000 ha)	Percentage (%)
High forest	872	34.7%	1 166	34.7%
Coppice forest	1 206	48.0%	1 612	48.0%
Coppice forest with standards	434	17.3%	581	17.3%
<b>Total</b>	<b>2 512</b>	<b>100.0%</b>	<b>3 359</b>	<b>100.0%</b>

Source: a) Distribution of Forests in Greece 1964, General Secretariat of Forests and Natural Environment (GSF&NE), Ministry of Agriculture  
b) First National Inventory of Forests 1992, GSF&NE, Ministry of Agriculture

### Images




Coppice Oak forest in Northern Greece

## Typology


<b>Simple coppice</b>	Used to be applied to all broadleaved species in the past. Today almost all of the beech has been converted and much of the oak coppice forests are being converted into high forests by extending the rotation and altering the method of stand tending. All evergreen broadleaved species forests (maquis) are managed as simple coppice.
<b>Coppice with standards</b>	Used to be applied traditionally in oak and chestnut coppice forests but in the nineties and after the chestnut blight infestation it was prohibited to manage chestnut coppice forests with standards in order to prevent the expansion of the disease. As an alternative it was suggested to extend the rotation time or to leave standards in groups, not individual trees. Coppice with standards management is applied regularly in all oak coppice forests of the country.
<b>Pollarding</b>	Practised locally in rather limited cases; not practised systematically. One exception is the pollarding of white mulberry trees for feeding the silkworm in sericulture or silk farming, but only on privately-owned mulberry plantations. Some livestock growers in rural Greece also occasionally use tree pollarding for animal fodder, but it is rather extensive and cannot be considered typical.
<b>Short rotation coppice</b>	Short rotation coppice is not officially applied in Greece. There is no law or other legal document for this particular management method specifically for energy purposes. One small exception of short rotation coppice concerns the basket willow ( <i>Salix viminalis</i> ), which was the raw material for the traditional basket making but today it is limited in very few places and the production is very small.
<b>Other types</b>	Coppice conversion into high forests: (1) coppice with standards can sometimes be implemented as an indirect method of conversion where a number of standard trees are retained individually or in groups at each rotation time and after several rotations the forest becomes uneven-aged and can be regenerated by seeds; (2) the extension of rotation time and stand tending by thinning is another indirect method of conversion.

## MAPS

Giorgos Mallinis and Gavriil Spyroglou



Overlaid map - range of the four main species that are coppiced in Greece (beech, sweet chestnut, oak and evergreen sclerophyllous shrubs)



Single maps - range of the four main species that are coppiced in Greece (beech, sweet chestnut, oak and evergreen sclerophyllous shrubs)

Data source: First National Inventory of Forests in Greece, 1992. GSF&NE, Ministry of Agriculture

## DESCRIPTION

### Gavriil Spyroglou

Coppice forests in Greece make up 65% of the forested area and 12% of the entire country (Ministry of Agriculture 1992). The main species are oaks (*Quercus* spp.) followed by chestnut (*Castanea sativa*), beech (*Fagus* spp.) and the evergreen broadleaves that make up the maquis. Other than chestnut, which can produce good quality wood in coppice rotations, the coppiced forests are characterized by very low growth rates, producing very low-value products such as firewood and charcoal. Most are grazed, either legally or illegally, and trees are still being pollarded by farmers and residents who keep a few domestic livestock animals. The aesthetic value is small because of

the large clear cut areas created by this management. As a result, many of these forests are not serving their required purpose, i.e. to provide an economic use (wood production), a protective function against soil erosion and aesthetic benefits. However, the great contribution of these forests is in mitigating climate change (Chatziphilippidis and Spyroglou 2004).

Coppice silviculture is a purely man-made management system that has been implemented in Europe since Roman times, based on the re-sprouting ability of broadleaf tree species. In the past, coppice management was the “child of necessity” and an easy management solution, but today it presents numerous ecological



and environmental problems which, in the context of sustainable, multifunctional, forest management should be directly addressed by a wide program of conversion to high forest. In Mediterranean environments, coppicing remains important because, despite the exhaustive logging, uncontrolled grazing and fires, intact ecosystems have been preserved in the coppice forests. Where forests are degraded, this is not necessarily linked to coppice management and the practice can contribute to improving both habitats and biodiversity with appropriate management. Other species, such as conifers or fast growing species, can co-exist in coppices, combining trees that regenerate from seed and those sprouting from coppice stools.

Conversion of coppice into high forests represents a change in management and can be achieved in two ways (Dafis 1966; Stamou 1981). Indirectly, by extending the rotation time so it equates to that of a high forest and managing the coppice stand as if it was of seedling origin. Alternatively, it can be achieved directly by changing the species, which usually takes place on very degraded sites and is achieved by planting conifers (pines). Coppice conversion in Greece has been going on for more than 90 years, with many fluctuations. The current coppice regime is based on the



Figure 1. Typical coppice forest in Taxiarchis, Chalkidiki

views of the 1950s and earlier. It is therefore appropriate to reconsider it under the current legislative framework and to develop a new strategic plan for a modern holistic approach that will meet today's challenges.

Mediterranean ecosystems in general, and coppice forests in particular, have been used over time for resources other than woody products. Non-timber forest products such as bark, forage, soil protection, mushrooms, fruits, honey and recreation are important. A critical evaluation of the whole spectrum of uses gives the real value of coppice forests. In this context, the Mediterranean coppice forests contribute to rural development, maintaining biodiversity and its associated economic values, ecosystem functions and services and last – but not least – are of considerable cultural importance.

## References

- Chatziphilippidis Gr. and Spyroglou G. 2004. Sustainable Management of coppice forests in Greece. In Anderson, F., Birot, Y., and Päävinen, R., (eds) *Towards the sustainable use of Europe's forests – forest ecosystem and landscape research: Scientific challenges and opportunities Proc 25-27 June 2003 Tours, France*. EFI Proceedings No 49, pp. 61 -70.
- Dafis, Sp. 1966. *Standorts-und Ertragskundliche Untersuchungen in Eichen-und Kastanien-niederwaeldern der N.O. Chalkidiki Thessaloniki*. [In Greek with German summary].
- Ministry of Agriculture 1992. *Results of the first National Forest Inventory*. Ministry of Agriculture, General secretariat of forests and natural environment, General directorate of development and protection of forests and natural environment, Athens, p. 134.
- Stamou, N. 1981. *Le Taillis simple de chênes en Grèce et ses traitements futurs, aspects économiques, conversion et enrésinement*. *Foret Méditerranée* III(2).

COST (European Cooperation in Science and Technology) is a funding agency for research and innovation networks. Our Actions help connect research initiatives across Europe and enable scientists to grow their ideas by sharing them with their peers. This boosts their research, career and innovation.

Published by:

Albert Ludwig University Freiburg  
Chair of Forest Utilization

Werthmannstr. 6  
D-79085 Freiburg  
Germany



[www.uni-freiburg.de](http://www.uni-freiburg.de)

**This article is part of the volume**

**“Coppice Forests in Europe”**

Printed by: Albert Ludwig University Freiburg Printing Press

Contact:

[www.eurocoppice.uni-freiburg.de](http://www.eurocoppice.uni-freiburg.de)  
[eurocoppice@fob.uni-freiburg.de](mailto:eurocoppice@fob.uni-freiburg.de)  
0049 (0)761 203 3789

### **Coppice Forests in Europe**

© 2018 Professur für Forstbenutzung, Albert-Ludwigs-Universität Freiburg, Freiburg i. Br., Germany

Editors: Alicia Unrau, Gero Becker, Raffaele Spinelli, Dagnija Lazdina, Natascia Magagnotti, Valeriu-Norocel Nicolescu, Peter Buckley, Debbie Bartlett and Pieter D. Kofman

ISBN 978-3-9817340-2-7

Recommended citations:

For the full volume: Unrau, A., Becker, G., Spinelli, R., Lazdina, D., Magagnotti, N., Nicolescu, V.N., Buckley, P., Bartlett, D., Kofman, P.D. (Eds.) (2018). *Coppice Forests in Europe*. Freiburg i. Br., Germany: Albert Ludwig University of Freiburg.

For individual chapters/articles: List of author(s) with surname(s) and initial(s). (2018). Chapter/article title. In A. Unrau, G. Becker, R. Spinelli, D. Lazdina, N. Magagnotti, V.N. Nicolescu, P. Buckley, D. Bartlett, P.D. Kofman (Eds.), *Coppice Forests in Europe* (pp. xx-xx). Freiburg i. Br., Germany: Albert Ludwig University of Freiburg.

The articles in this volume were developed within the context of COST Action FP1301 EuroCoppice (2013-2017). Numerous contributions were published as single, independent booklets during the course of the Action; they were subsequently reviewed and updated for this volume. A digital version of this volume, further results and more are available on the website: [www.eurocoppice.uni-freiburg.de](http://www.eurocoppice.uni-freiburg.de)

Design, layout & formatting: Alicia Unrau

Coppice image acknowledgements: Simple coppice (grey) based on a drawing by João Carvalho (pp. 46); Leaf vector originals designed by [www.freepik.com](http://www.freepik.com) (modified)

*Disclaimer: The views expressed in this publication are those of the authors and do not necessarily represent those of the COST Association or the Albert Ludwig University of Freiburg. Responsibility for content lies solely with the respective authors.*